

Superfluid Analog of the Davies-Unruh Effect

G. L. Comer

Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

(Dated: August, 1992)

Abstract

We produce an analog of the Davies-Unruh effect for superfluid helium four. There are two temperatures which result—one is associated with ordinary, or first, sound and the other with second sound.

Two of the most striking results of the investigations into quantum field theory on curved backgrounds are the Hawking effect [1] and the Davies-Unruh (D-U) effect [2, 3].¹ The Hawking effect predicts that black holes radiate particles thermally. The D-U effect predicts that a system interacting with some quantum field and following a particular non-inertial path undergoes transitions as if it were immersed in a thermal bath of excitations of the given quantum field, even though the quantum field itself is in a global Minkowski vacuum state.

It is no coincidence that both results have thermal attributes. In each case that which is responsible for the thermal effects is an event horizon.² In the black hole example the horizon is a feature of the geometric structure of the underlying spacetime; for the Davies-Unruh scenario a horizon exists because of the particular trajectory of the system. The main effect of a horizon is to delimit regions of spacetime between which information cannot be exchanged. In 1973, Bekenstein [6] pointed out, for the particular example of black holes, that this inability to exchange information can bring about a violation of the second law of thermodynamics. He argued that this violation can be prevented if black holes have, in addition to their usual physical attributes, thermal characteristics (i.e. entropy). The very general nature of Bekenstein’s reasoning suggests that it should always be the case that additional thermal properties result whenever an event horizon is present.

In 1981, Unruh [7] generalized the understanding behind these effects when he proposed an analog to the Hawking effect for irrotational perfect fluids. He considered an ordinary fluid which is steadily accelerated from zero velocity to supersonic flow. The place at which the transition to supersonic flow occurs, the sonic horizon, is the fluid analog of an event horizon. For example, the region of the fluid beyond the sonic horizon cannot communicate—through the exchange of fluid excitations (phonons)—with the starting place of the fluid flow (i.e. the fluid’s “asymptotic infinity” where its velocity is zero).³ Unruh found that the sonic horizon will emit phonons with a thermal distribution. The general-

¹ See also Fulling [4] for his preliminary contributions to the discovery of this effect.

² See Wald [5] for a suitable description of event horizons, in particular, the importance of causality (information cannot be exchanged instantaneously between two spatially separated points) in their definitions.

³ Here, to “communicate” means to send information *only* through the exchange of phonons. Of course one could penetrate this horizon with something like a photon, say, but this should not ruin the event horizon nature of the sonic horizon for the problem at hand since it is interactions with the fluid excitations that dominate the end results.

ization occurs because here it is causality as defined by the paths that phonons take, not light, which underpins the physics.

An important technical aspect of Unruh's derivation is that one can associate the fluid notions of sonic horizon and asymptotic infinity with an underlying "metric" in the mathematics describing the system. Unruh manipulates this metric in precisely the same way that real metrics are manipulated (e.g., arbitrary coordinate transformations, construction of metric compatible covariant derivatives, etc.) to obtain sonic horizon thermal radiation. It is the main purpose of this work to apply Unruh's technique in order to demonstrate an analog of the D-U effect for superfluid He⁴.

We use the two-fluid hydrodynamic model of superfluid He⁴ developed by Landau [8]. The two fluids are the so-called normal- and super-fluids. The equations of motion for this model are (see Putterman [9])

$$0 = \rho_{,t} + (\rho_n v_{nj} + \rho_s v_{sj})_{,j} , \quad (1)$$

$$0 = (\rho s)_{,t} + (\rho s v_{nj})_{,j} , \quad (2)$$

$$0 = v_{si,t} + v_{sj} v_{si,j} + \mu_{,i} , \quad (3)$$

$$0 = (\rho_n v_{ni} + \rho_s v_{si})_{,t} + (p \delta_{ij} + \rho_n v_{ni} v_{nj} + \rho_s v_{si} v_{sj})_{,j} , \quad (4)$$

where the repeated indices are summed over 1, 2, and 3, a subscript comma denotes partial differentiation, the subscripts n and s stand for normal- and super-fluid variables, respectively, any ρ , with a subscript or not, is a mass-density (ρ without a subscript is the total mass-density), v_j is the j^{th} component of a three-velocity, p is pressure, s is the specific entropy, μ is the chemical potential, and $\rho_s \equiv \rho - \rho_n$. These variables are all functions of the Newtonian time t and the spatial coordinates x_j . The eight equations of motion in Eqs. (1)–(4) are used to determine the eight independent degrees of freedom ρ , s , v_{ni} , and v_{si} . To have a well-posed problem we must also supplement these equations with equations of state. Defining $w_i \equiv v_{ni} - v_{si}$ and $w^2 \equiv w_j w_j$, then the equations of state have the functional dependence [9] $p = p[\rho, s, w^2]$, $\rho_n = \rho_n[\rho, s, w^2]$, and $\mu = \mu[\rho, s, w^2]$. If T is the temperature, then the thermodynamic variables must also satisfy [9]

$$dp = \rho d\mu + \rho s dT + \frac{1}{2} \rho_n dw^2 . \quad (5)$$

The form of Eq. (3) presupposes that the superfluid flow is irrotational, i.e. $\nabla \times \vec{v}_s = \vec{0}$.

This implies that there exists a scalar potential ϕ_1 such that

$$v_{si} = \phi_{1,i} . \quad (6)$$

As a consequence of Eq. (6), and an assumption about initial conditions, there is another vector field $A_i = (\rho_n w_i)/(\rho s)$ which is irrotational [9]. Thus there is a scalar potential ϕ_2 such that

$$A_i = \phi_{2,i} . \quad (7)$$

Because of the irrotational character of v_{si} and A_i we really have only four independent degrees of freedom; therefore, some of the above equations are redundant. By inserting Eq. (6) and Eq. (7) into Eqs. (1)–(4) we can reduce the number of equations of motion from eight to four. Inserting Eq. (6) into Eq. (3) yields

$$0 = \left(\phi_{1,t} + \frac{1}{2} \phi_{1,j} \phi_{1,j} + \mu \right)_{,i} \quad (8)$$

so that

$$0 = \phi_{1,t} + \frac{1}{2} \phi_{1,j} \phi_{1,j} + \mu + c_1(t) , \quad (9)$$

where $c_1(t)$ is an arbitrary function of t . Likewise, we can reduce Eq. (4) to find

$$0 = \phi_{2,t} + \frac{\rho s}{\rho_n} \phi_{2,j} \phi_{2,j} + \phi_{1,j} \phi_{2,j} + T + c_2(t) , \quad (10)$$

where $c_2(t)$ is another arbitrary function of t . Now we have as our independent degrees of freedom ϕ_1 , ϕ_2 , ρ , and s . Their equations of motion are Eqs. (1) and (2) (with Eqs. (6) and (7) inserted appropriately), Eq. (9), and Eq. (10), supplemented by the equations of state.

To demonstrate our analog of the D-U effect we use a linearized form of the equations of motion about some background solution. The variables ϕ_1 , ϕ_2 , ρ , s , p , T , μ , and ρ_n all take the form $\tilde{\mathcal{O}} = \mathcal{O} + \delta\mathcal{O}$, where $\tilde{\mathcal{O}}$ represents a full solution, $\mathcal{O}$ represents the background, and $\delta\mathcal{O}$ is some variation away from the background. The prefactor δ stands for a quantity which is “small” in the sense appropriate for this problem. We use a background which is homogeneous, isotropic, and static (i.e. all background functions ϕ_1 , ϕ_2 , etc. are independent of t and x_i). This implies that the thermodynamic variables p , T , μ , and ρ_n are functions of only ρ and s since the background velocities vanish.

Before going any further it is convenient to relate some of the thermodynamic derivatives which appear during the linearization with more familiar quantities, i.e. the heat capacities

at constant volume $C_V = T\partial s/\partial T|_\rho$ and constant pressure $C_p = T\partial s/\partial T|_p$ as well as the speeds of first and second sounds $u_1^2 = \partial p/\partial \rho|_s$ and $u_2^2 = (Ts^2\rho_s)/(C_V\rho_n)$, respectively [9]. In terms of C_V , C_p , u_1 , and u_2 we find

$$\left.\frac{\partial p}{\partial s}\right|_\rho = -\frac{\rho}{s}w_1^2, \quad (11)$$

$$\left.\frac{\partial \mu}{\partial \rho}\right|_s = \frac{1}{\rho} [u_1^2 + w_1^2], \quad (12)$$

$$\left.\frac{\partial \mu}{\partial s}\right|_\rho = -\frac{1}{s} \left[w_1^2 + \frac{\rho_n}{\rho_s} u_2^2 \right], \quad (13)$$

where

$$w_2^2 \equiv \left(\frac{\rho_s}{\rho_n} - 1 \right) w_1^2 \equiv \left(\frac{\rho_s}{\rho_n} - 1 \right) u_1 u_2 \left(\frac{\rho_n}{\rho_s} \left[1 - \frac{C_V}{C_p} \right] \right)^{1/2}. \quad (14)$$

Since the background velocities vanish we do not require any specific formulas for the thermodynamic derivatives of ρ_n (these derivatives will not appear in the linearized equations).

Inserting all variables which take the form $\tilde{\mathcal{O}} = \mathcal{O} + \delta\mathcal{O}$ into our equations of motion, keeping only terms linear in δ , and using the various thermodynamic quantities identified above, we find

$$\delta\rho = -\frac{\rho}{u_1^2} \left[1 - \frac{\rho_s}{\rho_n} \frac{w_1^4}{u_1^2 w_2^2} \right]^{-1} \left[\delta\phi_{1,t} + s \left(1 + \frac{\rho_s}{\rho_n} \frac{w_1^2}{u_2^2} \right) \delta\phi_{2,t} \right], \quad (15)$$

$$\delta s = -\frac{s^2 \rho_s}{u_2^2 \rho_n} \left[1 - \frac{\rho_s}{\rho_n} \frac{w_1^4}{u_1^2 u_2^2} \right]^{-1} \left[\left(1 + \frac{w_1^2}{u_1^2} \right) \delta\phi_{2,t} + \frac{1}{s} \frac{w_1^2}{u_1^2} \delta\phi_{1,t} \right], \quad (16)$$

$$0 = -\frac{1}{u_1^2} \delta\phi_{1,tt} + \left[1 + \frac{w_1^2}{u_1^2} \right] \delta\phi_{1,jj} + s \left[1 - \frac{u_2^2}{u_1^2} - \frac{w_2^2}{u_1^2} \right] \delta\phi_{2,jj}, \quad (17)$$

$$0 = -\frac{1}{u_2^2} \delta\phi_{2,tt} + \left[1 - \frac{w_1^2}{u_2^2} \right] \delta\phi_{2,jj} - \frac{1}{s} \frac{w_1^2}{u_2^2} \delta\phi_{1,jj}. \quad (18)$$

We have done some further manipulations to arrive at Eqs. (15)–(18): The linearized form of Eqs. (9) and (10) did not involve any derivatives of $\delta\rho$ or δs so they could be solved in terms of derivatives of $\delta\phi_1$ and $\delta\phi_2$ by purely algebraic means.

We now show that some of the terms which appear in Eqs. (15)–(18) can be neglected for temperatures $T = 10^{-3}$ K. In this temperature range the thermodynamics is dominated by phonons, the roton contributions being negligible [10, 11], and $u_1 = \sqrt{3}u_2 = 2.4 \times 10^4$ cm/sec. Using the various equations of states for superfluid He⁴ (see [10, 11]) and some basic thermodynamic identities, it can be shown that $s = u_1^2 \rho_n / T \approx 10^4 T^3$ erg/(gm K⁴) and

$(1 - C_V/C_p)^{1/2} \approx 10^{-2} T^2 \text{ K}^{-2}$. We have assumed that ρ_s is constant (i.e. independent of T) so that $\partial\rho/\partial T|_p \approx \partial\rho_n/\partial T|_p$. Using these approximations we find that the terms w_1^2/u_1^2 , $w_1^2/(su_2^2)$, and $s[1 - (u_2^2 + w_2^2)/u_1^2]$ are all much less than one for $T < 10^{-3} \text{ K}$. Therefore, the equations for $\delta\phi_1$ and $\delta\phi_2$ decouple and take the following approximate form

$$0 = -\frac{1}{u_a^2} \delta\phi_{a,tt} + \delta\phi_{a,jj} , \quad (19)$$

where $a = 1$ or 2 is just a label, not a tensor index.

Now we formally identify a metric from Eq. (19). This metric is obviously that associated with a flat spacetime:

$$ds_a^2 = -u_a^2 dt^2 + dx_i dx_i . \quad (20)$$

Note that time intervals are related to spatial intervals in this “spacetime” by the appropriate speed of sound u_a . The important structure here is the sonic-cone given by the congruence of curves which satisfy $ds_a^2 = 0$. It is this structure which allows for the existence of a horizon in superfluid He^4 and underpins our Davies-Unruh analog. In contrast to real Minkowski spacetime, we see in this approximation to the dynamics of superfluid He^4 two sonic cones—one for first sound and another for second sound.⁴ It is a general feature of Eqs. (15)–(18) that our system has two characteristic velocities for propagating excitations (it is not just a result of the approximations which generated Eq. (19)) [8, 9, 10, 11].

To extract our analog of the D-U effect we will use the arguments developed by Unruh in Ref. [3]. They involve the construction of “particle” detection devices. These devices register excitations of the quantum field under consideration, for us $\delta\phi_a$. The detectors are to be accelerated, with constant proper acceleration a_a , along the z -axis. (In this context, “proper” quantities are formed with the “metric” in Eq. (20).) The worldlines traced out by the detectors have the following parametrization:

$$z^2 - u_a^2 t^2 = 4R^2 \quad , \quad u_a t = z \tanh(\tau/2) , \quad (21)$$

and $x = y = 0$. The parameters τ and R are the analogs of Rindler coordinates for the metric in Eq. (20) [12]. In order to model how a detector registers excitations consider it to be a particle in a potential well. We will, for simplicity, let the potential be generated by

⁴ We can anticipate similar results for third and fourth sound. Moreover, one can easily see that the Hawking effect analog will have more than one horizon (which will emit thermal excitations).

a box. The box must confine the particle and at the same time permit it to interact with the fluid. Let the interaction of the particle with $\delta\phi_a$ be via the potential

$$V_{int} = \varepsilon_a \delta\phi_a \Phi , \quad (22)$$

where Φ is the wave function of the particle and ε_a is some coupling constant. Note that our choice of V_{int} requires the particle to interact either with $\delta\phi_1$ or $\delta\phi_2$ but not both.⁵ Also, since the gradient of $\delta\phi_a$ is a perturbation of a velocity, and velocities change sign under time reversal, V_{int} is time antisymmetric. This is not a serious problem here since our main result will be seen to be independent of the particular form of the interaction potential (i.e. no dependence on the coupling constant ε_a). The overall behavior of the particle is governed by its Schrödinger wave equation

$$i \frac{\partial}{\partial \tau} \Phi = \frac{2a_a}{u_a} \frac{1}{2m} \left(\frac{\partial^2}{\partial h^2} + \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + m^2 \frac{u_a^2}{R_c} h + \varepsilon_a \delta\phi_a \right) \Phi , \quad (23)$$

where m is the mass of the particle, $h \approx (R^2 - R_c^2)/R_c$ is its proper distance from the center R_c of the box, $a_a = u_a^2/2R_c$ is its proper acceleration, and the time derivative is with respect to the proper time of the particle (hence, the reason for the extra factor $2a_a/u_a$). The term proportional to m^2 just represents the force experienced by the particle because it is accelerating.

Through some mechanism, which need not be discussed here, the detector is constrained to follow a $R = R_c = \text{const}$ trajectory in the $0 \leq z < \infty$ and $-\infty < t < \infty$ region of a $u_a t - z$ spacetime diagram. As can be seen from Eq. (21) this path is a hyperbola. The asymptotes of this hyperbolic path are generated by the sonic cone (i.e. the 45° lines in our spacetime diagram) going through the origin ($t = z = 0$). It is not very difficult to see that the “future” asymptote (i.e. the one with positive slope) does represent an event horizon since a fluid excitation produced “beyond” it can never be registered by our detector.

The point of this construction is to see how the detector responds when it follows the given non-inertial path. A good indicator of this response is the transition probability per unit proper time P_{aj} for the detector (i.e. “particle”) to be excited from its ground state

⁵ In superfluid He⁴ it is possible to excite first sound modes without exciting second sound modes and vice versa. The reason for this is second sound modes are essentially thermal waves while first sound modes are associated with density disturbances. This follows from Eqs. (15) and (16) since in the small T limit $\delta\rho \propto \delta\phi_{1,t}$ and $\delta s \propto \delta\phi_{2,t}$. Also see Ref. [9].

to some excited energy eigenstate with energy E_j . We will not present the details here, but rather we refer the reader to Unruh [3] since the calculation of P_{aj} follows exactly as presented there.⁶ The final result is

$$P_{aj} = \frac{1}{\exp(2\pi u_a [E_j - E_0]/a_a) - 1} F_{det} , \quad (24)$$

where E_0 is the ground state energy and F_{det} just depends on the characteristics of the detector (ε_a , etc.). Note that Eq. (24) is precisely the form expected of a detector immersed in a thermal bath of excitations with temperature T_a given by (restoring $\hbar$ and k_b)

$$T_a = \frac{1}{2\pi} \frac{\hbar}{k_b} \frac{a_a}{u_a} . \quad (25)$$

This is the desired result. (As claimed earlier, it is independent of ε_a .) For each speed of sound u_a there is an associated temperature (cf. footnote 4). Notice that this is the same formula obtained by Davies and Unruh except that the speed of sound u_a replaces the speed of light. It is, up to factors of order one, what one would obtain from a dimensional analysis, using only the relevant physical parameters of this problem. Putting in the values for $\hbar$, k_b , and u_a gives $T_a \approx 10^{-13} (a_a/g)$ K where $g \approx 10^3$ cm/sec².

That we obtain two different temperatures is a necessary consequence of the fact that there are two different sonic-cone structures in superfluid He⁴. This is a required consistency check of the principle underlying our D-U analog—two sonic-cone structures necessarily implies two types of event horizons, which, in turn, suggests two different thermal effects.

To arrive at Eq. (25) we have formally identified certain characteristics of superfluid He⁴ with an underlying metric present in the mathematics. We then formally used this metric in ways that are familiar from differential geometry and quantum field theory on curved backgrounds. At best, our derivation can only be considered heuristic since it is not obvious how to interpret the “proper time” $\tau u_a/2a_a$ appearing in Eq. (23). It becomes especially problematic if one imagines a non-relativistic ion being the “detector” described by Eq. (23). The accepted understanding is that it is only Newtonian time, like t in Eqs. (1)–(4), which is the ion’s proper time. If $\tau u_a/2a_a$ has any real meaning it must be statistical, in essence, and perhaps, as Unruh states for his Hawking effect analog, “... (have) no physical significance except for the small-amplitude sonic motions of the medium” [7]. Whatever

⁶ The important steps are those which produce Eqs. (3.1a) through (3.10) in Unruh [3].

the interpretation, if one accepts the premise that horizons and thermal effects necessarily occur together, then one has little recourse but to believe that T_a must have real, and not merely formal, significance. The problem, then, is to figure out how it can be detected.

I would like to thank J. D. Bekenstein for introducing me to Unruh's irrotational fluid calculations and for suggesting that I look at superfluids. His guidance and insight were crucial for the completion of this work. I also wish to thank T. Piran, C. A. Fuchs and M. Schiffer for their support and Amos Ori for useful discussions in which some of the conceptual difficulties of this work were clarified. This research was supported by a Golda Meir fellowship with the Hebrew University of Jerusalem, Israel.

-
- [1] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).
 - [2] P. C. W. Davies, J. Phys. **A8**, 609 (1975).
 - [3] W. G. Unruh, Phys. Rev. D **14**, 870 (1976).
 - [4] S. A. Fulling, Phys. Rev. D **7**, 2850 (1972).
 - [5] R. M. Wald, *General Relativity* (The University of Chicago Press, Chicago, 1984).
 - [6] J. D. Bekenstein, Phys. Rev. **D7**, 2333 (1973).
 - [7] W. G. Unruh, Phys. Rev. Lett. **46**, 1351 (1981).
 - [8] L. Landau, J. Phys. USSR **5**, 71 (1941).
 - [9] J. Putterman, *Superfluid Hydrodynamics* (North-Holland, Amsterdam, 1974).
 - [10] F. London, *Superfluids. Vol. I. Macroscopic Theory of Superconductivity* (Wiley, New York, 1950).
 - [11] F. London, *Superfluids. Vol. II. Macroscopic Theory of Superfluid Helium* (Wiley, New York, 1954).
 - [12] W. Rindler, Am. J. Phys. **34**, 1174 (1996).

Appendix: Note Added in 2005

The work presented here is unpublished, and not otherwise available. The original preprint is of some historic interest: it was one of the earliest articles in the modern, post-1990 revival of “analog models” for curved spacetime. Every attempt has been made to

maintain the original form of the preprint. One typographical error, in Eq. (4), has been fixed. The referencing and footnoting are also different, since a modern typesetting program has been used.

The author wishes to thank Matt Visser and Haret Rosu for citing this work on a few occasions, and thus preventing it from passing away like Lennon and McCartney's Eleanor Rigby. He also thanks several colleagues and collaborators for encouraging him to post it on the e-print archive.

The author's current address is Department of Physics, Saint Louis University, St. Louis, MO 63156-0907, USA.